

HOSTED BY



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

ScienceDirect

[www.elsevier.com/locate/ssci](http://www.elsevier.com/locate/ssci)

# Sleep apnea and oxygen saturation in adults at 2640 m above sea level <sup>☆</sup>

Maria Angelica Bazurto Zapata\*, Elida Dueñas Meza, Claudia Jaramillo, Dario Maldonado Gomez, Carlos Torres Duque

Sleep Laboratory, Fundación Neumológica Colombiana (FNC), Colombia

## ARTICLE INFO

### Article history:

Received 13 December 2013

Received in revised form

8 May 2014

Accepted 14 May 2014

Available online 16 September 2014

### Keywords:

Sleep apnea

High altitude

Oxygen saturation

## ABSTRACT

**Purpose:** To describe the SpO<sub>2</sub> in wakefulness, sleep and during the apnea–hypopnea in adults living in Bogotá, located at 2640 m above sea level.

**Methods:** Descriptive observational study in adults referred for polysomnogram (PSG). A normal Apnea hypopnea index (AHI) was defined as  $\leq 5$  and obstructive sleep apnea (OSA) was classified as mild (AHI 5–15), moderate (AHI 15–30), and severe (AHI  $> 30$ ). T-test or ANOVA test for SpO<sub>2</sub> differences between groups was used.

**Results:** 1799 patients, 33% women. 222 (12.8%) did not have OSA (normal IAH), 268 (14.9%) mild OSA, 315 (17.5%) moderate, and 993 (55.2%) severe. In all cases a low SpO<sub>2</sub> (SpO<sub>2</sub>  $< 90\%$ ) was found.

The SpO<sub>2</sub> was lower when the AHI was higher, in wakefulness, in non-REM and in REM ( $p < 0.001$ ). For all grades of severity, SpO<sub>2</sub> decreased significantly from wakefulness to non-REM sleep and to REM sleep ( $p < 0.001$ ). Patients with severe OSA had higher desaturation during wakefulness ( $85.2 \pm 6.6\%$ ), non-REM sleep ( $83.1 \pm 7.7\%$ ), REM sleep ( $78.8 \pm 10.2$ ), and during events ( $75.1 \pm 9.1\%$ ).

**Conclusions:** Patients with OSA at 2640 m have nocturnal desaturation lower than 88%, which decreases with higher severity of OSA. The clinical impact of sleep disorders at this point may be greater than at sea level and should be studied.

© 2014 Published by Elsevier B.V. on behalf of Brazilian Association of Sleep. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

## 1. Introduction

Obstructive sleep apnea (OSA) is a frequent condition [1], characterized by recurrent respiratory pauses which lead to sleep fragmentation and intermittent desaturation.

For the definition of hypopnea and consequently, for the diagnosis of OSA, oxygen desaturation is involved as one of

its components [2,3]. Specifically, it is a necessary flow reduction (ventilation) associated with a decrease of more than 3–4% of oxygen saturation, (SpO<sub>2</sub>).

Even though persistent and intermittent nocturnal desaturation had been related with the presence of some consequences of OSA, such as pulmonary hypertension and atherosclerosis [4], the role that desaturation has on the pathogenesis, development

<sup>☆</sup>This study's cost were funded by the Fundación Neumológica Colombiana.

\*Correspondence to: Carrera 13 B No. 161-85. Tel.: +57 7428900.

E-mail address: [mbazurto@neumologica.org](mailto:mbazurto@neumologica.org) (M.A. Bazurto Zapata).

Peer review under responsibility of Brazilian Association of Sleep.

and treatment of pulmonary hypertension in patients with SAHS, is still obscure [5–7].

In Bogotá, a city situated at a high altitude (2640 m above sea level), the normal SpO<sub>2</sub> during wakefulness is found between 90 and 92% [8–12]. Unlike the observed sea level, a decrease of 3% or higher from this point, which is the criterion to define the presence of hypopnea, necessarily implicates the appearance of desaturation and its possible consequences. It is, therefore, very important to know the SpO<sub>2</sub> behavior during sleep at Bogotá's altitude as an initial step to investigate the role of altitude and the possible clinical repercussions of oxygen desaturation in patients with SAHS and if these findings justify a redefinition of this syndrome at this altitude.

The objective of this study was to describe the oxygen saturation during wakefulness sleep and apneas-hypopneas in adults at an altitude of 2640 m.

## 2. Materials and methods

Patients of 18 years or more, sent for basal polysomnogram at the Laboratory of Sleep of Fundación Neumológica Colombiana in Bogotá, city situated 2640 m above sea level, between the years of January 2004 and December 2005, were included.

The study was approved by the Ethics Committee of the institution in 2006 and the patients approved the realization of the polysomnogram.

Nocturnal polysomnogram were performed for a minimum of 4 h, with Alice 3 and 4 from Respiromics®, following international recommendations: 3 electroencephalogram channels (C3M2, C4M1, O1M2), 2 electro-oculography channels and electromyography (mandible and legs), electrocardiogram, air flow by pressure cannula and thermistor, respiratory effort with thoracic and abdominal belts, finger pulse oxymetry, snoring and position. The SpO<sub>2</sub> was measured by MARS Model 2001 oxymeters with the following technical specifications: Measurement amplitude of 0–100%, precision: 70–100% ± 25 and a resolution of 1%. The value of oxygen saturation and the pulse frequency were determined by an average of a 2 s sample.

Manual scoring of the records was performed. The sleep stages were classified according to international criteria during the years of study (Reschthaffen and Kales) [13]. Apnea was defined as the complete cessation of air flow during 10 s or more and hypopnea as the reduction of, at least, 30% of the flow for 10 s, associated to an arousal or an oxygen desaturation of 3% or more. The apnea-hypopnea index (AHI) was defined as the total number of apneas and hypopneas per hour of sleep. The severity of OSA was determined according to AHI: Mild: 5–15/h; moderate: 16 to 30/h; and severe: more than 30/h [14]. Oxygen desaturation was defined by a SpO<sub>2</sub> lower than 90%.

Patients who received oxygen during the study, the studies for CPAP titration, and those with less than 4 h, were excluded.

We used averages and standard deviations for the quantitative variables and the proportions for qualitative variables. To establish differences between the SpO<sub>2</sub> during the different stages of sleep for each group according to AHI, we used the *t* test for independent samples and the ANOVA test. To establish the degree of statistical significance of the different proportions, we used the *chi* square test. We used Pearson's coefficient to establish the correlation between AHI with age

and IMC. We considered it significant when *p* < 0.05. We used the statistical programs SPSS 10.0.

## 3. Results

We included 1799 patients sent to the Laboratory of Sleep for the realization of nocturnal basal polysomnogram, 33% were women and 222 (12.8%) had normal AHI. In the group of sleep apnea, 268 (14.9%) patients presented mild OSA, 315 (17.5%) moderate OSA, and 993 (55.2%) severe OSA. Age and body mass index increased significantly with the grade of severity for OSA (*p* < 0.001) (Table 1). There was a significant correlation, even weak, between AHI and BMI (*r* = 0.325, *p* < 0.001) and between AHI and age (*r* = 0.167, *p* < 0.001) (Table 1).

The average SpO<sub>2</sub> was inferior to 90% (oxygen desaturation) in all cases, during wakefulness or sleep. All subjects with normal AHI as well as all grades of severity of OSA, were found with a progressive and significant decrease in the average SpO<sub>2</sub>, when going from wakefulness to non-REM sleep and from non-REM sleep to REM sleep (*p* < 0.001). The higher AHI and OSA severity were found with less SpO<sub>2</sub>, during vigil as well as during non-REM and REM sleep (*p* < 0.001). Desaturation was larger (lower SpO<sub>2</sub>) during apnea or hypopnea events for all the severity groups of OSA. Average SpO<sub>2</sub> during the events were equally lower with the increased severity of OSA (*p* < 0.001) (Table 2 and Fig. 1).

Compared with the patients with mild SAHS, the patients with severe OSA had higher desaturation during wakefulness (87.5 ± 5.0 vs. 85.2 ± 6.6), non-REM sleeps (86.6 ± 5.4 vs. 83.1 ± 7.7), REM sleep (84.4 ± 8.7 vs. 78.8 ± 10.2) and during apneas and hypopneas (82 ± 6.9 vs. 75.1 ± 9.1) (*p* < 0.001).

## 4. Discussion

The main finding of this study was that patients with OSA at an altitude of 2640 m above sea level showed a significant oxygen desaturation during sleep, which decreases even more with the increased severity of OSA especially during REM sleep and during the events of apnea and hypopnea, reaching values as low as 78% during REM sleep and 75% during the above mentioned events.

The worst desaturation during REM sleep in patients with sleep apnea has been described and has been related with the duration of apneas during this sleep stage [15].

**Table 1 – Characteristics of the patients and distribution of the severity of SAHS.**

	N (%)	Age, years	BMI
Normal AHI	222 (12.8)	43.4 ± 16.0	26.0 ± 5.5
Mild OSA	268 (14.9)	50.2 ± 13.3	28.7 ± 5.6
Moderate OSA	315 (17.5)	53.1 ± 13.4	28.0 ± 5.7
Severe OSA	993 (55.2)	54.7 ± 12.8	31.1 ± 6.2

AHI: Apnea hypopnea Index (apnea hypopnea per hour of sleep).  
OSA: obstructive sleep apnea.  
BMI: body mass index.  
Values are averages ± SD or N(%).

**Table 2 – Average oxygen saturation during wakefulness, sleep stages and respiratory events.**

IAH	SpO <sub>2</sub> average wakefulness	SpO <sub>2</sub> average, during sleep			p*
		Non-REM	REM	Events	
Normal	88.7 ± 6.2	88.1 ± 5.7	87.0 ± 7.3	–	<0.001
Mild OSA	87.5 ± 5.0	86.6 ± 5.4	84.3 ± 8.7	82 ± 6.9	<0.001
Moderate OSA	86.4 ± 6.3	85.2 ± 6.4	83 ± 8.0	80 ± 8.0	<0.001
Severe OSA	85.2 ± 6.6	83.1 ± 7.7	78.8 ± 10.2	75.1 ± 9.1	<0.001
p**	<0.001	<0.001	<0.001	<0.001	


AHI: apnea hypopnea index (apneas-hypopneas per hour of sleep).

OSA: obstructive sleep apnea.

Values as average ± SD.

\* Value of p when comparing SpO<sub>2</sub> during the different stages of sleep in each group according to IAH.

\*\* Value of p when comparing SpO<sub>2</sub> of the groups according to IAH in each of the stages of sleep.



**Fig. 1 – SpO<sub>2</sub> (%) in wakefulness, during sleep and the respiratory events according to AHI. SpO<sub>2</sub>: oxygen saturation per oxymetry; AHI: Apnea hypopnea index (apneas-hypopneas per hour of sleep).**

Patz et al., have studied the effect of altitude upon AHI and did not find important differences between the minimal saturation in three different altitudes (0 m, 1370 m and 2423 m) [16].

Some studies have described the behavior of oxygen saturation during sleep. Gries et al. found that the average saturation in 350 normal subjects during sleep, in Cleveland, Ohio (173 m above sea level), was of 96.5%. The saturation decreased mildly with age, ranging from 96.8% in the group of 1–10 years of age, to 95.1% in the group of 60 years of age [17].

With altitude, the saturation values differ, however, the majority of the studies has been performed at more than 3000 m. In six normal subjects, the average SaO<sub>2</sub> during sleep were of 97.3%, 83.0% and 71.0% measured, respectively at 500 m, 4200 m, and 6400 m [18]. In another study, the saturation at extreme altitude (8400 m), was of 59% [19]. The studies during sleep in intermediate altitudes are few and were performed at simulated altitudes (normobaric hypoxia) [20,21].

Findings of the majority of studies performed at altitude are not comparable with the ones of our study, since they were performed after acute exposition to altitude, which can induce the appearance of central apneas and reflect the changes that are

detected with an early adaptation [22]. Rey de Castro et al., did respiratory polygraphy to truck drivers who worked at 2020 m and found out that 12% had central apnea and 10% had OSA. In the study group, the majority had intermittent exposition to the altitude. All desaturation indexes measured on the study were lower in central apnea when compared with OSA [23].

In Bogota (2640 m), the studies of normal values of PaO<sub>2</sub> have been conducted during wakefulness and involves normal and young subjects [8–12]. At this altitude, the average normal values of PaO<sub>2</sub>, for the range of age of our patients, are of around 65 mmHg, between 25 and 30% lower than at sea level.

However, at this altitude, PaO<sub>2</sub> is 60 mmHg, from which the curve of hemoglobin dissociation inclines and small changes of PaO<sub>2</sub> have significant repercussions on SpO<sub>2</sub>. All the patients with OSA had an average SpO<sub>2</sub> at desaturation levels (SpO<sub>2</sub> < 90%) with significant decreases according to the increased severity of OSA and going from wakefulness to non-REM sleep, from non-REM sleep to REM sleep and during events, reaching levels of SpO<sub>2</sub> inferior to 80% in patients with severe OSA during REM sleep and during events.

The repercussion of intermittent and persistent hypoxia, reflected upon the levels of desaturation in the patients of the study, might be very significant.

There are several ways to express the severity of desaturation during sleep such as average desaturation, the minimum, and the percentage of sleep time with saturation lower than 90%, all of these are comparable [19]. In this study, we have described the average oxygen saturation.

The strengths of this study were the significant number of patients included and that this is as our best knowledge, the first article describing the behavior of oxygen saturation during sleep in individuals with OSA, living at high altitude (above 2500 m), contributing for better understanding of this pathology among our population and in populations living in similar altitudes.

This study has some limitations: firstly, the population of the study was sent for a polysomnogram evaluation with a suspicion of a sleep disorder, for which there is no well defined normal group. From the total patients, only 12% had a normal AHI (< 5/h) with saturation during wakefulness of less than 90%, which suggests that some of these patients might have some pulmonary pathology. Additionally, we have not analyzed separately the population with diagnostic of hypoventilation, obesity, obstructive pulmonary disease, nor cardiovascular comorbidities.

Due to the fact that patients with more severe AHI were older and had higher BMI, additional future studies are dully required to evaluate the impact of these variables upon SpO<sub>2</sub> at Bogota's altitude, as well as the possible differences that might exist between genders.

In conclusion, in this group of patients, the oxygen saturation during sleep in patients with OSA at Bogota's altitude, is lower than 88%, decreasing progressively with the increased severity of OSA, especially during REM sleep. The impact of this upon the development of complications such as pulmonary hypertension in patients with SAHS at this time, is still undetermined.

### Conflict of interest

There are no conflicts of interests for any of the authors.

### REFERENCES

- [1] Young T, Peppard PE, Gottlieb DJ. Epidemiology of obstructive sleep apnea: a population health perspective. *Am J Respir Crit Care Med* 2002;165:1217-31.
- [2] Sleep-Related Breathing Disorders in Adults: recommendations for syndrome definition and measurement technics in clinical research. *SLEEP*, vol. 22; 1999. p. 667-87.
- [3] The AASM Manual for the Scoring of Sleep and Associated Events; 2007.
- [4] Savransky V, Nanayakkara A, Li J, Bevans S, Smith PL, Rodriguez A, et al. Chronic intermittent hypoxia induces atherosclerosis. *Am J Respir Crit Care Med* 2007;175:1290-7.
- [5] Chaouat A, Wetzenblum E. Pulmonary hemodynamics in the obstructive sleep apnea syndrome. *Chest* 1996;109:380-6.
- [6] Wetzenblum E, Krieger J. Daytime pulmonary hypertension in patients with obstructive sleep apnea. *Am Rev Resp Dis* 1988;138:345-9.
- [7] Laks L, Lehrhaft B, Grunstein RR, Sullivan CE. Pulmonary hipertensión in obstructive sleep apnea. *Eur Respir J* 1995;8:537-41.
- [8] Restrepo J, Reyes P, Vásquez P, Ardila M, Díaz-Granados B. Gasimetría arterial y alveolar en adultos sanos a nivel de Bogotá. *Acta Med Colomb* 1982;7(6):461-6.
- [9] Acevedo L E, Solarte I. Gasimetría arterial en adultos jóvenes a nivel de Bogotá. *Acta Med Colomb* 1982;9(1):7-14.
- [10] Duran M, Grandas N, Reyes P. Gasimetría arterial en adultos jóvenes sanos en Bogotá. *Rev Colomb Neumol* 1993;5(2):73-7.
- [11] Roa J, de Arroyo S, Maldonado D. Gasimetría arterial a diferentes edades en Bogotá. *Gasimetría arterial y alveolar en adultos sanos a nivel de Bogotá. Acta Med Colomb* 1988(13 suppl.): (451A(C 217).
- [12] Jaramillo M, Berbesí F. Valores normales de los gases arteriales y la brecha aniónica a nivel de Bogotá. Estudio en una muestra significativa. *Acta Med Colomb* 1988(13 suppl.):416 (C 221).
- [13] Rechtschaffen A, Kales A. A manual of Standardized Terminology, Techniques, and Scoring System for Sleep Stages of Human Subjects. Los Angeles, CA: Brain Information Service/Brain Research Institute; 1968.
- [14] Sleep-Related Breathing Disorders in Adults-AASM Task Force. *Sleep*, vol. 22, No. 5; 1999. p. 667-87.
- [15] Findley LJ, Wilhoit SC, Suratt PM. Apnea duration and hypoxemia during REM sleep in patients with obstructive sleep apnea. *Chest* 1985;87:432-6.
- [16] Patz D, Spoon M, et al. The effect of altitude descent on obstructive sleep apnea. *Chest* 2006;130:1744-50.
- [17] Gries RE, Brooks LJ. Normal oxyhemoglobin saturation during sleep: how low does it go? *Chest* 1996;110:1489-92.
- [18] Netzer NC, Strohl KP. Sleep and breathing in recreational climbers at an altitude of 4200 and 6400 m: observational study of sleep and patterning of respiration during sleep in a group of recreational climbers. *Sleep Breath* 1999;3(3):75-82.
- [19] Chaudhary B, Dasti S, et al. Hour-to-hour variability of oxygen saturation in sleep apnea. *Chest* 1998;113:719-22.
- [20] Hoshikawa M, Uchida S, Sugo T, Kumai Y, Hanai Y, Kawahara T. Changes in sleep quality of athletes under normobaric hypoxia equivalent to 2000-m altitude: a polysomnographic study. *J Appl Physiol* 2007;103:2005-11.
- [21] Kinsman T, Hahn A, Gore C, Wilshire B, Martin D, Chow CM. Respiratory events and periodic breathing in cyclists sleeping at 2650-m simulated altitude. *J Appl Physiol* 2002;92:2114-8.
- [22] Grocott MP, Martin D, Levett D, McMorrow R, Windsor J, Montgomery HE. Caudwell xtreme everest research group. Arterial blood gases and oxygen content in climbers on mount everest. *N Engl J Med* 2009;360:140-9.
- [23] Rey de Castro J, Ortiz O, Rosales-Mayor E, Ferreyra J, Liendo A, Liendo CH. Prevalence of sleep-disordered breathing in truck drivers at a mine located at high altitude. *Sleep Sci* 2013;6 (2):59-65.